

Chen's inequality in Lagrangian case

Teodor Oprea

University of Bucharest,

Faculty of Mathematics and Informatics

14 Academiei St., code 010014, Bucharest, Romania

e-mail: teodoroprea@yahoo.com

Abstract

In the theory of submanifolds, the following problem is fundamental: *to establish simple relationships between the main intrinsic invariants and the main extrinsic invariants of the submanifolds*. The basic relationships discovered until now [1, 2, 3, 4] are inequalities. To analyze these problems, we follow the idea of C. Udriște [6, 7, 8] that the method of constrained extremum is a natural way to prove geometric inequalities. We improve the Chen's inequality which characterizes a totally real submanifold of a complex space form. For that we suppose that the submanifold is Lagrangian and we formulate and analyze a suitable constrained extremum problem.

2000 *Mathematics subject classification*: 53C21, 53C24, 53C25, 49K35.

Keywords: constrained maximum, Chen's inequality, Lagrangian submanifolds.

1. Optimizations on Riemannian submanifolds

Let $(N, \tilde{g})$ be a Riemannian manifold of dimension m , M be a Riemannian submanifold of it, g be the metric induced on M by $\tilde{g}$ and $f : N \rightarrow R$ be a differentiable function.

In [5] we considered the constrained extremum problem

$$(1) \min_{x \in M} f(x).$$

The first result is

THEOREM 1. *If $x_0 \in M$ is the solution of the problem (1), then*

- i) $(\text{grad } f)(x_0) \in T_{x_0}^\perp M$,
- ii) *the bilinear form*

$$\alpha : T_{x_0}M \times T_{x_0}M \rightarrow R,$$

$$\alpha(X, Y) = \text{Hess}_f(X, Y) + \tilde{g}(h(X, Y), (\text{grad } f)(x_0)),$$

is positive semidefinite, where h is the second fundamental form of the submanifold M in N .

We shall use this theorem in order to find a inequality verified by the Chen's invariant of a Lagrangian submanifold in a complex space form.

2. Estimation of Chen's invariant of a Lagrangian submanifold in a complex space form

Let (M, g) be a Riemannian manifold of dimension n , and x a point in M . We consider the orthonormal frame $\{e_1, e_2, \dots, e_n\}$ in $T_x M$. The scalar curvature at x is defined by

$$\tau = \sum_{1 \leq i < j \leq n} R(e_i, e_j, e_i, e_j),$$

where R is the Riemann curvature tensor of (M, g) .

We denote $\delta_M = \tau - \min(k)$, where k is the sectional curvature at the point x . The invariant δ_M is called the Chen's invariant of Riemannian manifold (M, g) .

Let $(\tilde{M}, \tilde{g}, J)$ be a Kähler manifold of real dimension $2m$. A submanifold M of dimension n of $(\tilde{M}, \tilde{g}, J)$ is called a totally real submanifold if for any point x in M the relation $J(T_x M) \subset T_x^\perp M$ holds.

If, in addition, $n = m$, then M is called Lagrangian submanifold. For a Lagrangian submanifold, the relation $J(T_x M) = T_x^\perp M$ occurs.

A Kähler manifold with constant holomorphic sectional curvature is called a complex space form and is denoted by $\tilde{M}(c)$. The Riemann curvature tensor $\tilde{R}$ of $\tilde{M}(c)$ satisfies the relation

$$\tilde{R}(X, Y)Z = \frac{c}{4} \{ \tilde{g}(Y, Z)X - \tilde{g}(X, Z)Y + \tilde{g}(JY, Z)JX - \tilde{g}(JX, Z)JY + 2\tilde{g}(X, JY)JZ \}.$$

A totally real submanifold of real dimension n in a complex space form $\tilde{M}(c)$ of real dimension $2m$ verifies a Chen's inequality

$$\delta_M \leq \frac{n-2}{2} \left\{ \frac{n^2}{n-1} \|H\|^2 + (n+1) \frac{c}{4} \right\},$$

where H is the mean curvature vector of the Riemannian submanifold M of $\tilde{M}(c)$.

REMARK. i) If M is a totally real submanifold of real dimension n in a complex space form $\tilde{M}(c)$ of real dimension $2m$, then

$$A_{JY}X = -Jh(X, Y) = A_{JX}Y, \forall X, Y \in \mathcal{X}(M).$$

ii) Let $m = n$ (M is Lagrangian in $\widetilde{M}(c)$). If we consider the point $x \in M$, the orthonormal frames $\{e_1, \dots, e_n\}$ in $T_x M$ and $\{Je_1, \dots, Je_n\}$ in $T_x^\perp M$, then

$$h_{jk}^i = h_{ik}^j, \forall i, j, k \in \overline{1, n},$$

where h_{jk}^i is the component after Je_i of the vector $h(e_j, e_k)$.

With these ingredients we prove the next result which can be regarded as an obstruction to Lagrangian isometric immersions of a Riemannian manifold into a complex space form.

THEOREM 2. *Let M be a Lagrangian submanifold in complex space form $\widetilde{M}(c)$ of real dimension $2n$, $n \geq 3$. Then*

$$\delta_M \leq \frac{(n-2)(n+1)}{2} \frac{c}{4} + \frac{n^2}{2} \frac{2n-3}{2n+3} \|H\|^2.$$

PROOF. We consider the point $x \in M$, the orthonormal frames $\{e_1, \dots, e_n\}$ in $T_x M$ and $\{Je_1, \dots, Je_n\}$ in $T_x^\perp M$, $\{e_1, e_2\}$ being an orthonormal frame in the 2-plane which minimize the sectional curvature at the point x in $T_x M$.

By using Gauss's equation $\widetilde{R}(X, Y, Z, U) = \widetilde{R}(X, Y, Z, U) - g(h(X, Z), h(Y, U)) + g(h(X, U), h(Y, Z)), \forall X, Y, Z, U \in \mathcal{X}(M)$ and the fact that $\widetilde{M}(c)$ is a complex space form, we obtain

$$(1) \quad \tau = \frac{n(n-1)}{2} \frac{c}{4} + \sum_{r=1}^n \sum_{1 \leq i < j \leq n} h_{ii}^r h_{jj}^r - \sum_{r=1}^n \sum_{1 \leq i < j \leq n} (h_{ij}^r)^2,$$

$$(2) \quad R(e_1, e_2, e_1, e_2) = \frac{c}{4} + \sum_{r=1}^n h_{11}^r h_{22}^r - \sum_{r=1}^n (h_{12}^r)^2.$$

By subtracting the relations (1) and (2), we find

$$(3) \quad \delta_M = \frac{(n-2)(n+1)}{2} \frac{c}{4} + \sum_{r=1}^n \left(\sum_{3 \leq j \leq n} (h_{11}^r + h_{22}^r) h_{jj}^r + \sum_{3 \leq i < j \leq n} h_{ii}^r h_{jj}^r - \sum_{3 \leq j \leq n} (h_{1j}^r)^2 - \sum_{2 \leq i < j \leq n} (h_{ij}^r)^2 \right).$$

By using the symmetry in the three indexes of h_{ij}^k , we can write

$$\begin{aligned} (4) \quad \delta_M &\leq \frac{(n-2)(n+1)}{2} \frac{c}{4} + \sum_{r=1}^n \left(\sum_{3 \leq j \leq n} (h_{11}^r + h_{22}^r) h_{jj}^r + \sum_{3 \leq i < j \leq n} h_{ii}^r h_{jj}^r \right) - \\ &- \sum_{3 \leq j \leq n} (h_{1j}^1)^2 - \sum_{3 \leq j \leq n} (h_{1j}^j)^2 - \sum_{2 \leq i < j \leq n} (h_{ij}^i)^2 - \sum_{2 \leq i < j \leq n} (h_{ij}^j)^2 = \\ &= \frac{(n-2)(n+1)}{2} \frac{c}{4} + \sum_{r=1}^n \left(\sum_{3 \leq j \leq n} (h_{11}^r + h_{22}^r) h_{jj}^r + \sum_{3 \leq i < j \leq n} h_{ii}^r h_{jj}^r \right) - \\ &- \sum_{3 \leq j \leq n} (h_{11}^j)^2 - \sum_{3 \leq j \leq n} (h_{jj}^1)^2 - \sum_{2 \leq i < j \leq n} (h_{ii}^j)^2 - \sum_{2 \leq i < j \leq n} (h_{jj}^i)^2 = \end{aligned}$$

$$\begin{aligned}
&= \frac{(n-2)(n+1)}{2} \frac{c}{4} + \sum_{r=1}^n \left(\sum_{3 \leq j \leq n} (h_{11}^r + h_{22}^r) h_{jj}^r + \sum_{3 \leq i < j \leq n} h_{ii}^r h_{jj}^r \right) - \\
&- \sum_{3 \leq j \leq n} (h_{11}^j)^2 - \sum_{3 \leq j \leq n} (h_{jj}^1)^2 - \sum_{\substack{i,j \in \overline{2,n} \\ i \neq j}} (h_{jj}^i)^2.
\end{aligned}$$

Let us consider the quadratic forms $f_1, f_2, f_r : R^n \rightarrow R, r \in \overline{3,n}$ defined respectively by

$$\begin{aligned}
f_1(h_{11}^1, h_{22}^1, \dots, h_{nn}^1) &= \sum_{3 \leq j \leq n} (h_{11}^1 + h_{22}^1) h_{jj}^1 + \sum_{3 \leq i < j \leq n} h_{ii}^1 h_{jj}^1 - \sum_{3 \leq j \leq n} (h_{jj}^1)^2, \\
f_2(h_{11}^2, h_{22}^2, \dots, h_{nn}^2) &= \sum_{3 \leq j \leq n} (h_{11}^2 + h_{22}^2) h_{jj}^2 + \sum_{3 \leq i < j \leq n} h_{ii}^2 h_{jj}^2 - \sum_{3 \leq j \leq n} (h_{jj}^2)^2, \\
f_r(h_{11}^r, h_{22}^r, \dots, h_{nn}^r) &= \sum_{3 \leq j \leq n} (h_{11}^r + h_{22}^r) h_{jj}^r + \sum_{3 \leq i < j \leq n} h_{ii}^r h_{jj}^r - \\
&- (h_{11}^r)^2 - \sum_{\substack{j \in \overline{2,n} \\ j \neq r}} (h_{jj}^r)^2.
\end{aligned}$$

We need the maximum of f_1 and f_3 . For f_2 and $f_r, r \in \overline{4,n}$, we can solve similar problems.

We start with the problem

$$\max f_1,$$

$$\text{subject to } P : h_{11}^1 + h_{22}^1 + \dots + h_{nn}^1 = k^1,$$

where k^1 is a real constant.

The first three partial derivatives of the function f_1 are

$$\begin{aligned}
(5) \quad \frac{\partial f_1}{\partial h_{11}^1} &= \sum_{3 \leq j \leq n} h_{jj}^1, \\
(6) \quad \frac{\partial f_1}{\partial h_{22}^1} &= \sum_{3 \leq j \leq n} h_{jj}^1, \\
(7) \quad \frac{\partial f_1}{\partial h_{33}^1} &= h_{11}^1 + h_{22}^1 + \sum_{4 \leq j \leq n} h_{jj}^1 - 2h_{33}^1.
\end{aligned}$$

As for a solution $(h_{11}^1, h_{22}^1, \dots, h_{nn}^1)$ of the problem in question, the vector $(\text{grad})(f_1)$ is normal at P , from (5), (6) and (7) we obtain

$$(8) \quad h_{11}^1 + h_{22}^1 = 3h_{jj}^1 = 3a^1, \forall j \in \overline{3,n}.$$

By using the relation $h_{11}^1 + h_{22}^1 + h_{33}^1 + \dots + h_{nn}^1 = k^1$, from (8) we obtain $3a^1 + (n-2)a^1 = k^1$. Consequently

$$(9) \quad a^1 = \frac{k^1}{n+1}.$$

As f_1 is obtained from the function studied in Chen's inequality (see [5]) by subtracting some square terms, $f_1|_P$ will have the Hessian seminegative definite. Consequently the point $(h_{11}^1, h_{22}^1, \dots, h_{nn}^1)$ given by the relations (8) and (9) is a global maximum point, and hence

$$(10) \quad f_1 \leq 3a^1(n-2)a^1 + C_{n-2}^2(a^1)^2 - (n-2)(a^1)^2 = \frac{(a^1)^2}{2}(n+1)(n-2).$$

From (9) and (10), it follows

$$(11) \quad f_1 \leq \frac{(k^1)^2}{2} \frac{n-2}{n+1} = \frac{(n)^2}{2} \frac{n-2}{n+1} (H^1)^2.$$

Similarly we obtain

$$(12) \quad f_2 \leq \frac{(n)^2}{2} \frac{n-2}{n+1} (H^2)^2.$$

Further on, we shall consider the problem

$$\max f_3,$$

$$\text{subject to } P : h_{11}^3 + h_{22}^3 + \dots + h_{nn}^3 = k^3,$$

where k^3 is a real constant.

The first four partial derivatives of the function f_3 are

$$(13) \quad \frac{\partial f_3}{\partial h_{11}^3} = \sum_{3 \leq j \leq n} h_{jj}^3 - 2h_{11}^3,$$

$$(14) \quad \frac{\partial f_3}{\partial h_{22}^3} = \sum_{3 \leq j \leq n} h_{jj}^3 - 2h_{22}^3,$$

$$(15) \quad \frac{\partial f_3}{\partial h_{33}^3} = h_{11}^3 + h_{22}^3 + \sum_{4 \leq j \leq n} h_{jj}^3,$$

$$(16) \quad \frac{\partial f_3}{\partial h_{44}^3} = h_{11}^3 + h_{22}^3 + \sum_{\substack{3 \leq j \leq n \\ j \neq 4}} h_{jj}^3 - 2h_{44}^3.$$

For a solution $(h_{11}^1, h_{22}^1, \dots, h_{nn}^1)$ of the problem in question, the vector $(\text{grad})(f_3)$ is colinear to $(1, 1, \dots, 1)$.

By using (13), (14), (15) and (16) we obtain

$$(17) \quad h_{11}^3 = h_{22}^3 = 3a^3,$$

$$(18) \quad h_{33}^3 = 12a^3,$$

$$(19) \quad h_{jj}^3 = 4a^3, \forall j \in \overline{4, n}.$$

As $h_{11}^3 + h_{22}^3 + h_{33}^3 + \dots + h_{nn}^3 = k^3$, from (17), (18) and (19), one gets

$$(20) \quad a^3 = \frac{k^3}{4n+6}.$$

With an argument similar to those in the previous problem we obtain that the point $(h_{11}^3, h_{22}^3, \dots, h_{nn}^3)$ given by the relations (17), (18), (19) and (20) is a global maximum point. Therefore

$$(21) \quad f_3 \leq 6a^3 12a^3 + 6a^3(n-3)4a^3 + C_{n-3}^2 16(a^3)^2 -$$

$$-18(a^3)^2 - (n-3)16(a^3)^2 = 2(a^3)^2(2n-3)(2n+3).$$

From (20) and (21) we obtain $f_3 \leq \frac{(k^3)^2}{2} \frac{2n-3}{2n+3} = \frac{n^2}{2} \frac{2n-3}{2n+3} (H^3)^2$.

Similarly one gets

$$(22) \quad f_r \leq \frac{n^2}{2} \frac{2n-3}{2n+3} (H^r)^2, \forall r \in \overline{3, n}.$$

As $\frac{n-2}{n+1} < \frac{2n-3}{2n+3}$, $\forall n \geq 3$, from (11), (12) and (22), it follows

$$(23) \quad f_r \leq \frac{n^2}{2} \frac{2n-3}{2n+3} (H^r)^2, \forall r \in \overline{1, n}.$$

By using the relations (4) and (23) we have

$$(24) \quad \delta_M \leq \frac{(n-2)(n+1)}{2} \frac{c}{4} + \frac{n^2}{2} \frac{2n-3}{2n+3} \sum_{r=1}^n (H^r)^2 = \frac{(n-2)(n+1)}{2} \frac{c}{4} + \frac{n^2}{2} \frac{2n-3}{2n+3} \|H\|^2.$$

REMARK. B.Y. Chen, F. Dillen, L. Verstraelen, L. Vrancken showed in [4] that an Lagrangian submanifold, of real dimension $2n$, $n \geq 3$, of a complex space form $\widetilde{M}(c)$, satisfying the equality

$$(25) \delta_M = \frac{n-2}{2} \left\{ \frac{n^2}{n-1} \|H\|^2 + (n+1) \frac{c}{4} \right\},$$

is minimal.

Now, this result is an immediate consequence of the previous inequality.

From (24) and (25), it follows $\frac{n-2}{2} \frac{n^2}{n-1} \|H\|^2 \leq \frac{n^2}{2} \frac{2n-3}{2n+3} \|H\|^2$, whence $\frac{n^2}{2} \|H\|^2 \left(\frac{2n-3}{2n+3} - \frac{n-2}{n-1} \right) \geq 0$.
As $\frac{2n-3}{2n+3} - \frac{n-2}{n-1} < 0$, $\forall n \geq 3$, we infer that $\|H\| = 0$, so M is a minimal submanifold in $\widetilde{M}(c)$.

Acknowledgment. I would like to thank to Professor C. Udriște, who has always been generous with his time and advice.

References

- [1] B.Y. Chen, 'Some pinching classification theorems for minimal submanifolds', *Arch. Math.* **60** (1993), 568-578.
- [2] B.Y. Chen, 'Some new obstructions to minimal Lagrangian isometric immersions', *Japan. J. Math.* **26** (2000), 105-127.
- [3] B.Y. Chen, 'Ideal Lagrangian immersions in complex space forms', *Math. Proc. Cambridge Philos. Soc.* **128** (2000), 511-533.
- [4] B.Y. Chen, F. Dillen, L. Verstraelen, L. Vrancken, 'Totally real submanifolds of CP^n satisfying a basic equality', *Arch. Math.* **63** (1994), 553-564.
- [5] T. Oprea, 'Optimizations on Riemannian Submanifolds', *Analele Univ. Buc.*, **LIV** 1 (2005), 127-136.
- [6] C. Udriște, E. Tănăsescu, 'Minime și maxime ale funcțiilor reale de variabile reale', Editura Tehnică, București, 1980.
- [7] C. Udriște, 'Convex functions and optimization methods on Riemannian manifolds', Kluwer Academic Publishers Group, Dordrecht, 1994.
- [8] C. Udriște, O. Dogaru, I. Țevy, 'Extrema with nonholonomic constraints', Geometry Balkan Press, Bucharest 2002.